

Analysis of $B \rightarrow K_0^*(1430)\ell^+\ell^-$ Decay in the Supersymmetric Models

V. Bashiry^{1*}, M. Bayar^{2†}, K. Azizi^{2‡},

¹ Engineering Faculty, Cyprus International University,
Via Mersin 10, Turkey

² Physics Department, Middle East Technical University,
06531 Ankara, Turkey

Abstract

In this paper, we study the branching ratios of $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay, where $K_0^*(1430)$ being the p-wave scalar meson, in the supersymmetric models. The branching ratios are estimated when short and long distance effects are taken into account. It is found that the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ ($\ell = \mu, \tau$) decay is measurable at LHC. Measurement of these branching ratios for the semileptonic rare $B \rightarrow K_0^*(1430)\ell^+\ell^-$, in particular, at low q^2 region can give valuable information about the nature of interactions within Standard Model or beyond.

*e-mail: bashiry@ciu.edu.tr

†e-mail: mbayar@metu.edu.tr

‡e-mail: e146342@metu.edu.tr

1 Introduction

The Standard Model (SM) is in perfect agreement with all confirmed collider data, but requires a missing ingredient. The SM is not regarded as a full theory, since it can not address some issues i.e., gauge and fermion mass hierarchy, matter- antimatter asymmetry, number of generations, the nature of the dark matter, the unification of fundamental forces and so on. For these reasons, the SM can be considered as an effective theory of some fundamental theory at low energy.

Supersymmetry (SUSY) is regarded as the most plausible extension of the SM in order to shed light on some of the issues mentioned above [1]. It is an essential ingredient in string theory and the most-favoured candidate for unifying the all known interactions including gravity. It would help stabilize the hierarchy of mass scales between m_W and the Planck mass, by canceling the quadratic divergences in the radiative corrections to the mass-squared of the Higgs boson [2].

Two types of study can be conducted to explore supersymmetric particles (sparticles). In the direct search, the center of mass energy of colliding particles must be increased to produce SUSY particles at the TeV scale, and hence be accessible to the LHC. On the other hand, we can investigate SUSY effects, indirectly. The sparticles can contribute to the quantum loop level. As a result, flavor changing neutral current (FCNC) transition induced by quantum loop level can be considered as a good tool for studying the possible effects of sparticles (there are many studies in this regard, for the most recent studies see Ref. [3] and the references therein).

The FCNC processes induced by $b \rightarrow s(d)$ transitions provide the most sensitive and stringiest test for the SM at one loop level, since they are forbidden in SM at tree level [4, 5]. Despite smallness of the branching ratios of FCNC decays, quite intriguing results are obtained in ongoing experiments. The inclusive $B \rightarrow X_s \ell^+ \ell^-$ decay is observed in BaBar [6] and Belle collaborations. These collaborations also measured exclusive modes $B \rightarrow K \ell^+ \ell^-$ [7–9] and $B \rightarrow K^* \ell^+ \ell^-$ [10]. The experimental results on these decays are in a good agreement with theoretical estimations [11–13] the results of which can be used to constrain new physics (NP) effects.

There is another class of rare decays induced by $b \rightarrow s$ transition, such as $B \rightarrow K_{02}^*(1430) \ell^+ \ell^-$ in which B meson decays into p-wave scalar meson. The decays $B \rightarrow K_2^*(1430) \ell^+ \ell^-$ and $B \rightarrow K_0^*(1430) \ell^+ \ell^-$ are studied in [14, 15]. The main problem in these studies is the calculation of the transition form factors. Transition form factors of these decays in the framework of light front quark model [16] and 3-point QCD sum rules are estimated in [17], [18] and [15], respectively.

In the present work we investigate the possible effects of sparticles on the branching ratio of $B \rightarrow K_0^*(1430) \ell^+ \ell^-$ decay.

The paper is organized as follows: In section 2, we calculate the decay amplitude of the $B \rightarrow K_0^*(1430) \ell^+ \ell^-$ decay within SUSY models. Section 3 is devoted to the numerical analysis and discussion of the considered decay and our conclusions.

2 Decay amplitude of the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay in the SUSY models

The exclusive $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay is described at quark level by $b \rightarrow s\ell^+\ell^-$ transition. The effective Hamiltonian responsible for the $b \rightarrow s\ell^+\ell^-$ transition in SUSY models (see, for example, Ref. [19]) is:

$$\mathcal{H}_{eff} = \frac{G_F \alpha V_{tb} V_{ts}^*}{2\sqrt{2}\pi} \left[C_9^{eff}(m_b) \bar{s} \gamma_\mu (1 - \gamma_5) b \bar{\ell} \gamma^\mu \ell + C_{10}(m_b) \bar{s} \gamma_\mu (1 - \gamma_5) b \bar{\ell} \gamma^\mu \gamma_5 \ell \right. \\ \left. - 2m_b C_7(m_b) \frac{1}{q^2} \bar{s} i \sigma_{\mu\nu} q^\nu (1 + \gamma_5) b \bar{\ell} \gamma^\mu \ell + C_{Q_1} \bar{s} (1 + \gamma_5) b \bar{\ell} \ell + C_{Q_2} \bar{s} (1 + \gamma_5) b \bar{\ell} \gamma_5 \ell \right], \quad (1)$$

The terms proportional to the Wilson Coefficients C_{Q_1} and C_{Q_2} come from the new Higgs bosons (NHB) exchange diagrams exist in supersymmetric theories. The Wilson coefficients C_7 , C_9^{eff} and C_{10} are already exist in the SM. $C_9^{eff}(\hat{s}) = C_9 + Y(\hat{s})$, where $Y(\hat{s}) = Y_{\text{pert}}(\hat{s}) + Y_{\text{LD}}$ contains both the perturbative part $Y_{\text{pert}}(\hat{s})$ and long-distance part $Y_{\text{LD}}(\hat{s})$ (see Ref. [11]). The explicit expressions of C_7 , C_9^{per} and C_{10} in the SM can be found in [4]. Y_{LD} is usually parameterized by using Breit–Wigner ansatz,

$$Y_{\text{LD}} = \frac{3\pi}{\alpha^2} C^{(0)} \sum_{V_i = \psi(1s) \dots \psi(6s)} \alpha_i \frac{\Gamma(V_i \rightarrow \ell^+ \ell^-) m_{V_i}}{m_{V_i}^2 - q^2 - i m_{V_i} \Gamma_{V_i}},$$

where α is the fine structure constant and $C^{(0)} = 0.362$.

The phenomenological factors α_i for the $B \rightarrow K(K^*)\ell^+\ell^-$ decay can be determined from the condition that they should reproduce correct branching ratio relation

$$\mathcal{B}(B \rightarrow J/\psi K(K^*) \rightarrow K(K^*)\ell^+\ell^-) = \mathcal{B}(B \rightarrow J/\psi K(K^*)) \mathcal{B}(J/\psi \rightarrow \ell^+\ell^-),$$

where the right-hand side is determined from experiments. Using the experimental values of the branching ratios for the $B \rightarrow V_i K(K^*)$ and $V_i \rightarrow \ell^+\ell^-$ decays, for the lowest two J/ψ and ψ' resonances, the factor α_i takes the values: $\alpha_1 = 2.7$, $\alpha_2 = 3.51$ (for K meson), and $\alpha_1 = 1.65$, $\alpha_2 = 2.36$ (for K^* meson). The values of α_i used for higher resonances are usually the average of the values obtained for the J/ψ and ψ' resonances. In order to determine the branching ratio for the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay with the inclusion of long distance effects, the values of α_i are borrowed from $B \rightarrow K\ell^+\ell^-$ and $B \rightarrow K^*\ell^+\ell^-$ decays, namely, we choose $\alpha = 1$ and $\alpha = 2$ and performed numerical calculations with these values.

One has to sandwich Eq. (1) between initial meson state $B(p)$ and final meson state $K_0^*(1430)(p')$ in order to obtain the amplitude for the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay. Thus, the matrix elements $\langle K_0^* | \bar{s} \gamma_\mu (1 - \gamma_5) b | B \rangle$ and $\langle K_0^* | \bar{s} i \sigma_{\mu\nu} q^\nu (1 + \gamma_5) b | B \rangle$ are needed. These matrix elements are parameterized in terms of the form factors as follows:

$$\langle K_0^*(1430)(p') | \bar{s} \gamma_\mu \gamma_5 b | B(p) \rangle = f_+(q^2) \mathcal{P}_\mu + f_-(q^2) q_\mu, \quad (2)$$

$$\langle K_0^*(1430)(p') | \bar{s} i \sigma_{\mu\nu} q^\nu \gamma_5 b | B(p) \rangle = \frac{f_T(q^2)}{m_B + m_{K_0^*}} [\mathcal{P}_\mu q^2 - (m_B^2 - m_{K_0^*}^2) q_\mu], \quad (3)$$

where $\mathcal{P}_\mu = (p + p')_\mu$ and $q_\mu = (p - p')_\mu$. By multiplying both sides of Eq. (2) with q^μ one can obtain the expression in terms of form factors for $\langle K_0^*(1430)(p') | \bar{s}\gamma_5 b | B(p) \rangle$.

$$\langle K_0^*(1430)(p') | \bar{s}\gamma_5 b | B(p) \rangle = -\frac{1}{m_b - m_s} [f_+(q^2)\mathcal{P} \cdot q + f_-(q^2)q^2], \quad (4)$$

Using above Hamiltonian and definitions of form factors, the decay amplitude for $B \rightarrow K_0^*\ell^+\ell^-$ can be written as follows:

$$\mathcal{M}(B \rightarrow K_0^*\ell^+\ell^-) = \frac{G_F\alpha V_{tb}V_{ts}^*}{2\sqrt{2}\pi} \left[-A_1 P_\mu \bar{\ell}\gamma^\mu \ell - A_2 P_\mu \bar{\ell}\gamma^\mu \gamma_5 \ell - A_3 \bar{\ell}\gamma_5 \ell - A_4 \bar{\ell}\ell \right], \quad (5)$$

where

$$\begin{aligned} A_1 &= C_9 f_+ + \frac{2m_b C_7 f_T}{m_B + m_{K_0^*}} \\ A_2 &= C_{10} f_+ \\ A_3 &= 2C_{10} m_\ell f_- + \frac{C_{Q_2}}{m_b - m_s} [(m_B^2 + m_{K_0^*}^2) f_+ + q^2 f_-] \\ A_4 &= \frac{C_{Q_1}}{m_b - m_s} [(m_B^2 + m_{K_0^*}^2) f_+ + q^2 f_-]. \end{aligned}$$

Using Eqs. (1)–(5), we get the following expression for the differential decay width:

$$\begin{aligned} \frac{d\Gamma}{dq^2} &= \frac{G_F^2 \alpha^2}{8192 m_B \pi^5} |V_{tb} V_{ts}^*|^2 v \sqrt{\lambda(1, r, \hat{s})} \left\{ \frac{4}{3} (|A_1|^2 + |A_2|^2) (-3 + v^2) \left[q^4 \right. \right. \\ &\quad - 2q^2 (m_B^2 + m_{K_0^*}^2) + (m_B^2 - m_{K_0^*}^2)^2 \Big] + 16 |A_2|^2 m_\ell^2 \left[q^2 - 2(m_B^2 + m_{K_0^*}^2) \right] \\ &\quad \left. \left. - 4q^2 |A_3|^2 + 4 |A_4|^2 (4m_\ell^2 - q^2) - 6m_\ell (A_2 A_3^* + A_2^* A_3) (m_B^2 - m_{K_0^*}^2) \right] \right\}, \quad (6) \end{aligned}$$

where $\hat{s} = \frac{q^2}{m_B^2}$, $v = \sqrt{1 - \frac{4m_\ell^2}{q^2}}$, $r = m_{K_0^*}^2/m_B^2$, and $\lambda(1, r, \hat{s}) = 1 + r^2 + \hat{s}^2 - 2\hat{s} - 2(1 + \hat{s})$.

3 Numerical results

In this section, we present the branching ratio for the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay for muon and tau leptons. The main input parameters are the form factors for which we use the results of three-point QCD sum rules [15].

The values of the form factors at $q^2 = 0$ are [15]

$$\begin{aligned} f_+(0) &= 0.31 \pm 0.08, \\ f_-(0) &= -0.31 \pm 0.07, \\ f_T(0) &= -0.26 \pm 0.07, \end{aligned} \quad (7)$$

where the errors are due to the variation of Borel parameters. From the sum rules expressions for the form factors it is argued that they are very sensitive to the magnitude of the leptonic decay constant $f_{K_0^*}$ of the $K_0^*(1430)$ meson [15].

$$\begin{aligned} f_+(0) &= 0.34 \pm 0.08 , \\ f_-(0) &= -0.34 \pm 0.07 , \\ f_T(0) &= -0.29 \pm 0.07 . \end{aligned} \tag{8}$$

The best fit for the q^2 dependence of the form factors can be written in the following form:

$$f_i(\hat{s}) = \frac{f_i(0)}{1 - a_i \hat{s} + b_i \hat{s}^2} , \tag{9}$$

where $i = +, -$ or T and $\hat{s} = q^2/m_B^2$. The values of the parameters $f_i(0)$, a_i and b_i are given in Table 1.

	$f_i(0)$	a_i	b_i
f_+	0.31 ± 0.08	0.81	-0.21
f_-	-0.31 ± 0.07	0.80	-0.36
f_T	-0.26 ± 0.07	0.41	-0.32

Table 1: Form factors for $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay in a three-parameter fit.

The kinematical interval of the dilepton invariant mass q^2 is $4m_\ell^2 \leq q^2 \leq (m_B - m_{K_0^*})^2$ in which the long distance effects (the charmonium resonances) can give substantial contribution. The dominant contribution to the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decays comes from the two low lying resonances J/ψ and ψ' , in the interval of $8 \text{ GeV}^2 \leq q^2 \leq 14 \text{ GeV}^2$. In order to minimize the hadronic uncertainties we discard this subinterval by dividing the kinematical region of q^2 for muon:

$$\begin{aligned} \text{I} \quad & 4m_\ell^2 \leq q^2 \leq (m_{J\psi} - 0.02 \text{ GeV})^2 , \\ \text{II} \quad & (m_{J\psi} + 0.02 \text{ GeV})^2 \leq q^2 \leq (m_{\psi'} - 0.02 \text{ GeV})^2 , \\ \text{III} \quad & (m_{\psi'} + 0.02 \text{ GeV})^2 \leq q^2 \leq (m_B - m_{K_0^*})^2 . \end{aligned}$$

and for tau:

$$\begin{aligned} \text{I} \quad & 4m_\ell^2 \leq q^2 \leq (m_\psi - 0.02 \text{ GeV})^2 , \\ \text{II} \quad & (m_\psi + 0.02 \text{ GeV})^2 \leq q^2 \leq (m_B - m_{K_0^*})^2 . \end{aligned}$$

In Fig. (1) and (2) we present the dependence of the differential branching ratio for the $B \rightarrow K_0^*(1430)\ell^+\ell^-$ where $\ell = \mu, \tau$ decay on q^2 , as well as its dependence on q^2 due solely

Wilson Coefficients	C_7^{eff}	C_9	C_{10}
SM	-0.313	4.334	-4.669
SUSY I	+0.3756	4.7674	-3.7354
SUSY II	+0.3756	4.7674	-3.7354
SUSY III	-0.3756	4.7674	-3.7354

Table 2: Wilson Coefficients in SM and different SUSY models but without NHBs contributions.

Wilson Coefficients	C_{Q_1}	C_{Q_2}
SM	0	0
SUSY I	0	0
SUSY II	6.5 (16.5)	-6.5 (-16.5)
SUSY III	1.2 (4.5)	-1.2 (-4.5)

Table 3: Wilson coefficient corresponding to NHBs contributions within SUSY I, II and III models [19]. The values in the bracket are for the τ .

to short distance effects ($\alpha = 0$ case). We also use the results of Ref. [19] for the values of Wilson coefficients in the supersymmetric theories (Table 2 and 3).

Taking into account the q^2 dependence of the form factors given in Eq. (9), performing integration over q^2 , and using the total life time $\tau_B = 1.53 \times 10^{-12} s$ [20], we get the following results for the branching ratios when only short distance contribution is taken into account:

$$\mathcal{B}(B \rightarrow K_0^*(1430)\mu^+\mu^-) = \begin{cases} 1.05 \times 10^{-7} & \text{SUSY I ,} \\ 2.08 \times 10^{-7} & \text{SUSY II ,} \\ 1.10 \times 10^{-7} & \text{SUSY III ,} \end{cases}$$

$$\mathcal{B}(B \rightarrow K_0^*(1430)\tau^+\tau^-) = \begin{cases} 9.54 \times 10^{-10} & \text{SUSY I ,} \\ 1.25 \times 10^{-8} & \text{SUSY II ,} \\ 2.69 \times 10^{-9} & \text{SUSY III .} \end{cases}$$

Taking long distance effects into account in the above-mentioned kinematical regions, we get the following branching ratios for muon:

$$\mathcal{B}(B \rightarrow K_0^*(1430)\mu^+\mu^-) = \begin{cases} 1.05 \times 10^{-7} & \text{region I ,} \\ 8.98 \times 10^{-9} & \text{region II ,} \\ 1.56 \times 10^{-10} & \text{region III ,} \end{cases} \quad \text{for SUSY I,}$$

$$\mathcal{B}(B \rightarrow K_0^*(1430)\mu^+\mu^-) = \begin{cases} 1.73 \times 10^{-7} & \text{region I ,} \\ 3.71 \times 10^{-8} & \text{region II ,} \\ 3.25 \times 10^{-9} & \text{region III ,} \end{cases} \quad \text{for SUSY II,}$$

and

$$\mathcal{B}(B \rightarrow K_0^*(1430)\mu^+\mu^-) = \begin{cases} 1.08 \times 10^{-7} & \text{region I ,} \\ 1.02 \times 10^{-8} & \text{region II ,} \\ 2.83 \times 10^{-10} & \text{region III ,} \end{cases} \quad \text{for SUSY III.}$$

and for tau:

$$\mathcal{B}(B \rightarrow K_0^*(1430)\tau^+\tau^-) = \begin{cases} 5.77 \times 10^{-10} & \text{region I ,} \\ 3.43 \times 10^{-10} & \text{region II ,} \end{cases} \quad \text{for SUSY I,}$$

$$\mathcal{B}(B \rightarrow K_0^*(1430)\tau^+\tau^-) = \begin{cases} 4.67 \times 10^{-9} & \text{region I ,} \\ 5.84 \times 10^{-9} & \text{region II ,} \end{cases} \quad \text{for SUSY II,}$$

and

$$\mathcal{B}(B \rightarrow K_0^*(1430)\tau^+\tau^-) = \begin{cases} 1.21 \times 10^{-9} & \text{region I ,} \\ 1.15 \times 10^{-9} & \text{region II ,} \end{cases} \quad \text{for SUSY III.}$$

at $f_{K_0^*} = 340 \text{ MeV}$.

It follows from these results that the dominant contribution comes from term proportional to C_7 in region I (low invariant mass region), and this can be attributed to the existence of the factor $1/q^2$. At LHCb 10^{11} – 10^{12} pairs are expected to be produced, the expected number of events for the $B \rightarrow K_0^*(1430)\mu^+\mu^-$ decay in the low invariant mass region is of the order of 10^4 – 10^5 . Since, this region is sensitive to the sign of C_7 in the SUSY I model, the study of branching ratio in this region can give valuable information about the SUSY effects. Moreover, comparing the value of the branching ratio for the $B \rightarrow K_0^*(1430)\mu^+\mu^-$ decay with and without long distance effects can give valuable results to check structure of the effective Hamiltonian. The smallness of the value of $\mathcal{B}(B \rightarrow K_0^*(1430)\tau^+\tau^-)$ can be attributed to the small phase volume of this decay. Furthermore, SUSY models can enhance the branching ratio up to one order of magnitude with respect to the SM values for both μ and τ cases. The significant discrepancy in the non-resonance regions (low q^2 and high q^2 regions) can be studied for the effects of not only NHBs but also for NP effects.

To sum up, we study the semileptonic rare $B \rightarrow K_0^*(1430)\ell^+\ell^-$ decay in the supersymmetric theories. We show that the branching ratio is very sensitive to the SUSY parameters.

The branching ratio is enhanced up to one order of magnitude with respect to the corresponding SM values. The results of which can be used to indirect search for the SUSY effects in future planned experiments at LHC.

Note added

After this work had been completed, we noticed recent paper [21] in which $B \rightarrow K_0^*(1430)\ell^+\ell^-$ were studied in supersymmetric theories. Their numerical analysis based on the light cone QCD sum rules results for the values of form factors. In our study we consider three-point QCD sum rules results for the values of form factors and the branching ratio is different from those presented in Ref. [21].

Acknowledgments

The authors thank T. M. Aliev for his useful discussions.

References

- [1] J. Ellis, arXiv:0902.0357
- [2] E. Witten, Phys. Lett. B **105** 267 (1981).
- [3] R. N. Hodgkinson, AIP Conf. Proc. **1078**, 375 (2009) [arXiv:0810.0034 [hep-ph]].
- [4] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [5] A. Ali, Int. J. Mod. Phys. A **20**, 5080 (2005).
- [6] B. Aubert *et. al*, BaBar Collaboration, Phys. Rev. Lett. **93**, 081802 (2004).
- [7] M. I. Iwasaki *et. al*, BELLE Collaboration, Phys. Rev. D **72**, 092005 (2005).
- [8] K. Abe *et. al*, BELLE Collaboration, Phys. Rev. Lett. **88**, 021801 (2002).
- [9] B. Aubert *et. al*, BaBar Collaboration, Phys. Rev. Lett. **91**, 221802 (2003).
- [10] A. Ishikawa *et. al*, BELLE Collaboration, Phys. Rev. Lett. **91**, 261601 (2003).
- [11] P. Colangelo, F. De Fazio, P. Santorelli and E. Scrimieri, Phys. Rev. D **53**, 3672 (1996); Errata D **57**, 3186 (1998); A. Ali, P. Ball, L. T. Handoko and G. Hiller, Phys. Rev. D **61**, 074024 (2000); A. Ali, E. Lunghi, C. Greub and G. Hiller, Phys. Rev. D **66**, 034002 (2002).
- [12] T. M. Aliev, H. Koru, A. Özpıneci, M. Savcı, Phys. Lett. B **400**, 194 (1997); T. M. Aliev, A. Özpıneci, M. Savcı, Phys. Rev. D **56**, 4260 (1997); D. Melikhov, N. Nikitin and S. Simula, Phys. Rev. D **57**, 6814 (1998).
- [13] G. Burdman, Phys. Rev. D **52**, 6400 (1995); J. L. Hewett and J. D. Walls, Phys. Rev. D **55**, 5549 (1997); C. H. Chen and C. Q. Geng, Phys. Rev. D **63**, 114025 (2001).
- [14] S. Rai Choudhury, A. S. Cornell, G. C. Joshi and B. H. J. McKellar, Phys. Rev. D **74**, 054031 (2006).
- [15] T. M. Aliev, K. Azizi and M. Savcı, Phys. Rev. D **76**, 074017 (2007) [arXiv:0710.1508 [hep-ph]].
- [16] H. Y. Cheng, C. K. Chua, and C. W. Huang, Phys. Rev. D **69**, 074025 (2004).
- [17] H. Y. Cheng, C. K. Chua, Phys. Rev. D **69**, 094007 (2004).
- [18] C. H. Chen, C. Q. Geng, C. C. Lih and C. C. Liu, Phys. Rev. D **75**, 074010 (2007).
- [19] M. J. Aslam, Y. M. Wang and C. D. Lu, Phys. Rev. D **78**, 114032 (2008) [arXiv:0808.2113 [hep-ph]].
- [20] W. M. Yao *et. al*, Particle Data Group, J. Phys. G **33**, 1 (2006).
- [21] M. J. Aslam, C. D. Lu and Y. M. Wang, arXiv:0902.0432 [hep-ph].




Figure 1: The dependence of the $d\Gamma/dq^2$ on q^2 for $B \rightarrow K_0^* \mu^+ \mu^-$.




Figure 2: The dependence of the $d\Gamma/dq^2$ on q^2 for $B \rightarrow K_0^* \tau^+ \tau^-$.